

0830 0059

The Goldacres Mines Co. owns 14 claims 29 miles by road southwest of Esowawo, in the low hills at the east front of the Shoshone Range, near the common corner of T. 27-28 N., R. 46-47 E. The property was worked for gold in the period 1934-41, and was equipped with a camp and 50-ton cyanide mill. Scheelite was found in 1943 in small amounts that proved non-commercial.

The tungsten mineralization is southwest of the gold ore body in irregular bodies of tascite formed in a thick limestone sequence. The tascite is cut by quartz-calcite veins that carry small amounts

584

of scheelite, galena, pyrite, and chalcopyrite. The size and shape of the tascite bodies was not readily determinable in 1943 because the outcrops and other exposures were so limited in extent. Much of the exposed tascite is barren or contains less than 0.1 percent of WO_3 . In a shallow trench at the south end of the property, a layer 2 feet wide exposed for a length of 15 feet is estimated to contain 1.0 percent of WO_3 . At another prospect, a 75-foot crosscut driven from the bottom of a 100-foot shaft exposed a quartz-calcite-sulfide vein 4 to 18 inches wide. This vein contains pockets rich in scheelite; a grab sample from a sorted ore pile assayed 1.25 percent of WO_3 .

According to Vanderburg, a little scheelite accompanied gold

Vanderburg, W. O., Reconnaissance of mining districts in Lander
County, Nevada: U. S. Bur. Mines Inf. Cir. 7043, pp. 43-44,
1939.

in placer deposits dredged by the Mill Gulch Placer Mining Co. from
1937 to 1939. The deposits are in Mill Gulch near Tenabo, 24 miles
southwest of Beowawe, on the east slope of the Shoshone Range.