1370 0024

- a. Ferguson (Delamar) district, Lincoln County, Nevada.
- b. Geographic coordinates: 37°24' N., 114°49' W.

c. Discovered in 1892 and produced through 1909. Recorded production 1902-32: 317,100 oz. Ag, some copper and lead.

d. References: Hewett and others, 1936.

e. Adequacy of our present knowledge: ?

Topographic coverage: ?

All-2 g. Major mineralogic and geologic features: Disseminated deposits and veinlets in shattered Paleozoic quartzite.

Silver in the United States

(Data sheets for individual mining districts, prepared in conjunction with metallogenic map for 1960 International Geological Congress.)

Authorship:

E. T. McKnight - All districts west of the Mississippi River, except most of those silver-producing districts containing less than 1,000 tons of lead or zinc in the following states: Arizona, New Mexico, Nevada, Oregon and Washington. Also the following silver districts in 4 Ash Peak, of the states mentioned: Vulture, and Helvetia, Ariz.; Miami, Globe, Apache, Black Range, Chloride Flat, Georgetown and Lake Valley, New Mexico: Ashwood and Granite, oregon; Deertrail, Nespelan and Ruby-Conconully, Washington.

White Vine district, Michigan.

A. V. Heyl, Jr. - All districts east of the Mississippi River (except Whate Price,

Harry Klemic and W. L. Newman - Silver districts not associated with lead or zinc, in Arizona, New Mexico, Nevada, Oregon, and Washington (except as listed above).

Size categories of deposits (as penciled in left margins)

	0	1	/ 2	3
l.	Less than 1,000 tons	1,000 to 50,000 tons	50,000 to 1,000,000 tons	More than 1,000,000 tons
b	n	11	**	tr
n.	tt.	Ħ	12	n .
S	Less than 100,000 oz.	100,000 to 5,000,000 oz.	5,000,000 to 50,000,000 oz.	More than 1,000,000 oz.
u	Less than 10,000 oz.	10,000 to 100.000 oz.	100,000 to 1,000,000 oz.	More than 1,000,000 oz.

(NOTE: Categories for Au are less certain than for others.)

District No. on metallogenic map penciled at lower right.