1760 0020

- a. Mammoth (Ellsworth) district, Nye County, Nevada.
- b. Geographic coordinates: 38°58'N., 117°45'W.

Cu-0

c. Status of exploitation: Discovered at Ellsworth in 1863 by Indians.

86.0

d. References: Kral, 1951.

Zn-0

e. Adequacy of our present knowledge: ?

Ag-1

f. Topographic coverage: ?

g. Major mineralogic and geologic features: Veins in Tertiary rhyolite dikes cutting Triassic(?) metavolcanic rocks.

Silver in the United States

(Data sheets for individual mining districts, prepared in conjunction with metallogenic map for 1960 International Geological Congress.)

Authorship:

E. T. McKnight - All districts west of the Mississippi River, except most of those silver-producing districts containing less than 1,000 tons of lead or zinc in the following states: Arizona, New Mexico, Nevada, Oregon and Washington. Also the following silver districts in 4 Ash Peak of the states mentioned: Vulture and Helvetia, Ariz.; Miami, Globe, Apache, Black Range, Chloride Flat, Georgetown and Lake Valley, New Mexico: Ashwood and Granite, oregon; Deertrail, Nespelem and Ruby-Conconully, Washington.

White Vine district, Michigan.

A. V. Heyl, Jr. - All districts east of the Mississippi River (except Whate Pine,

Harry Klemic and W. L. Newman - Eilver districts not associated with lead or zinc, in Arizona, New Maxico, Nevaña, Oregon, and Washington (except as listed above).

Size categories of deposits (as penciled in left margins)

	0	1		2	3	
	Less than 1,000 tons	1,000 to 50,000 tons		50,000 to 00,000 tons	More than 1,000,000	tons
	11	Ħ		ŧ	11	
	11	f1		tf.	Ħ	
	Less than	100,000 to	į	,000,000 to	More than	
Œ	100,000 oz.	5,000,000 oz.	5	,000,000 oz.	₹,000,000	OZ.
u	Less than	10,000 to		100,000 to	More than	
	10,000 oz.	100.000 oz.	. j a	.,000,000 02.	1,000,000	oz.

(NOTE: Categories for Au are less certain than for others.)

District No. on metallogenic map peniled at lower right