

2550 0004

The Lincoln and Greenhorn property holdings consist of four Lincoln and 3 Greenhorn lode claims situated about $\frac{1}{2}$ mile south of Bartlett Peak, and 102 miles by improved, dirt roads, and trail northwest of Winnemucca. In 1954, the property was leased to the Golden Scheelite Mining and Milling Co.

The mine workings consist of 2 trenches, 20 and 60 feet in length, and a pit 50 feet long, 15 feet wide and 15 feet deep.

Rocks exposed in the mine area include quartz monzonite, tactite and biotite feldspar-quartz schist. Bartlett Peak, several hundred feet east is composed of granitic rock. The contact of the quartz monzonite and the metamorphosed sedimentary rocks strikes northeast and dips steeply southwest. Along this contact for a distance of 300 feet the sedimentary rocks have been altered to tactite in small isolated sections. Scheelite associated with powellite occurs in most of the tactite exposed.

A production of 6,5058 dry tons of ore containing 0.17 percent WO_3 was shipped to the Gatchell Mine Inc. Ore that ranges from 0.25 to 0.59 percent WO_3 remains in the mine workings. The property has been idle since 1954.

Osgood Range

Scheelite mineralization occurs in the northern half of the Osgood Range (K3) in tactite bodies formed by the metamorphism of limestone along the contacts of an intrusive granodiorite stock. Although the occurrence of scheelite in the area was known before 1917, very little tungsten was produced until the fall of 1942 when part of the Getchell mill was converted to treat tungsten ores. The tungsten unit was installed jointly by the Union Carbide Nuclear Co. and Getchell Mines, Ind.

Gold and tungsten are the only economically important metals in the district; although zinc, copper, silver, molybdenum, lead and bismuth occur in the tactite in small veins or as replacements in the metamorphosed rocks. Scheelite is the only tungsten mineral which has been recognized in the district, but most of the scheelite has a moderate content of powellite.

Limestone adjoins the igneous rock along about 1/3 of the margin of the stock, and tactite is present along 60 to 70 percent of the limestone-granodiorite contact. The tactite bodies vary greatly in size and in tungsten content. The ore bodies that have been mined ranged in grade from 0.34 to 0.8 percent WO_3 .

Mining operations on the 12 deposits that have been worked in the district have been mostly confined to surface pits and glory holes.

Production from the Riley mine in 1956, was from underground mining operations in open stopes.

The mines and prospects in the area are briefly described as follows: