S20, 38N, 42E

Valley View (Humboldt Co.) 136

Item 22

3680022

The Valley View mine, locally known as the Saunder's property consists of 3 and a fraction lode claims situated, about 23 miles northeast of Golconda and 5 miles south of the Getchell mine.

Rocks exposed in the area comprise limestone interbedded with thin sections of argillite which in this area lie adjacent to a granodiorite contact that trends north-south and dips 45° to 80° east. Scheelite mineralization is present in a narrow tactite zone on the limestone-granodiorite contact which on the surface can be traced for a distance of 2,200 feet. Metamorphism has to some extent altered the argillite to hornfels, but scheelite mineralization in sufficient amounts to form ore, occurs only in a 5 to 6 foot zone of tactite on the granodiorite contact. The tactite in this zone is composed chiefly of garnet and calcite, with minor amounts of quartz, molybdenite, pyrite and scheelite. The tactite in places is stained with copper carbonates. The scheelite content varies from 0.3 to 1.0 percent WO_3.

The principal development openings consist of a south adit 255 in length and a north adit 315 feet long from which several small stopes were opened. Numerous trenches, pits and cuts, partially expose the contact beyond the north adit workings.

Shipments of selectively mined ore from the north adit workings amounted to 1,530 tons that averaged 0.95 percent WO_3. USBM Unpubl. data, 1963

USBM unpublished rept. Humboldt Co., Valley View Sec. 20,T38N,R42E Owned by Don Degner

136

Item 22

The Valley View mine, locally known as the Saunder's property consists of 3 and a fraction lode claims situated, about 23 miles northeast of Golconda and 5 miles south of the Getchell mine.

Rocks exposed in the area comprise limestone interbedded with thin sections of argillite which in this area lie adjacent to a granodiorite contact that trends north-south and dips 45° to 80° east. Scheelite mineralization is present in a narrow tactite zone on the limestone-granodiorite contact which on the surface can be traced for a distance of 2,200 feet. Metamorphism has to some extent altered the argillite to hornfels, but scheelite mineralization in sufficient amounts to form ore, occurs only in a 5 to

6 foot zone of tactite on the granodiorite contact. The tactite in this zone is composed chiefly of garnet and calcite, with minor amounts of quartz, molybdenite, pyrite and scheelite. The tactite in places is stained with copper carbonates. The scheelite content varies from 0.3 to 1.0 percent WO_3.

The principal development openings consist of a south adit 255 in length and a north adit 315 feet long from which several small stopes were opened. Numerous trenches, pits and cuts, partially expose the contact beyond the north adit workings.

Shipments of selectively mined ore from the north adit workings amounted to 1,530 tons that averaged 0.95 percent WO_3.