

see silver peak district  
# 43900068

38600065

101

ITEM 76

BULLETIN OF THE GEOLOGICAL SOCIETY OF AMERICA

VOL. 20, PP. 223-264, PLS. 6-9

JULY 15, 1909

CONTRIBUTION TO THE GEOLOGY OF THE SILVER PEAK QUADRANGLE, NEVADA<sup>1</sup>

BY H. W. TURNER

(Presented before the Cordilleran Section of the Society December 31, 1908)

CONTENTS

	Page
Introduction .....	224
Geographic features .....	226
Extent and topography of the quadrangle.....	226
Drainage .....	227
Water supply .....	227
Precipitation .....	228
Scenery .....	228
Desert varnish .....	229
Pre-Cambrian complex .....	230
Location and composition of the complex.....	230
The gneisses .....	230
The schists .....	231
The white granite.....	232
Granitic quartz-veins .....	235
Diorite dikes .....	237
Paleozoic sedimentary series.....	238
Algonkian rocks .....	238
The Paleozoic era.....	238
Lower Cambrian .....	239
Upper Cambrian .....	242
The Ordovician sediments.....	243
Tertiary sedimentary series.....	243
Esmeralda formation .....	243
General character .....	243
Areal distribution of the beds.....	244
Basal conglomerate of the Esmeralda formation.....	245
Tertiary detrital-slope breccias.....	245
Conglomerate beds .....	246
Thickness of the beds.....	246
Neocene river gravels.....	246

<sup>1</sup> Published by permission of the Director of the U. S. Geological Survey. Manuscript received by the Secretary of the Society January 13, 1909.

KE NIPIGON BASIN  
oring upland, would place  
fect.  
eologic History?  
succeeded the Cretaceous  
opography had reached a  
nigon basin. There were  
e coastal plain sediments,  
Archean oldland and the  
r of different points in  
olten diabase were forced  
se parts of the district  
removed the diabase was  
masses or broke across the  
e of the laccolitic masses  
e edge of the escarpment  
lowland. Through other  
ks of the lowland, fluid  
wed each other with suffi-  
ount of liquid rock col-  
er cuesta, between it and  
that it overflowed upon  
eck cooled very slowly  
per part. Probably the  
y throughout the whole  
ent of the diabase-(pos-  
of the underlying sedi-  
he fossil post-Cretaceous  
cover. The post-diabase  
very similar to that of  
pouring of the diabase  
he region, disconnecting  
ts, modifying locally the  
y affecting the rates of  
colitic sills and capping  
ct than the underlying  
ne numerous mesas and

basins (except Superior) are