## 4790 0017

- a. Tempiute district, Lincoln County, Nevada.
- b. Geographic coordinates: 37°39' N., 115°37' W.
- c. Status of exploitation: Discovered in the sixties. Recorded production: 15,469 oz. Ag from 1924 to 1932. Since then about 150,000 oz. Ag was produced.
- d. References: Hewett and others, 1936
- e. Adequacy of our present knowledge: ?
- f. Topographic coverage: ?
- g. Major mineralogic and geologic features: Veins in lower Paleozoic limestone.

Ag-1

Au-l

## Silver in the United States

(Data sheets for individual mining districts, prepared in conjunction with metallogenic map for 1960 International Geological Congress.)

## Authorship:

E. T. McKnight - All districts west of the Mississippi River, except most of those silver-producing districts containing less than 1,000 tons of lead or zinc in the following states: Arizona, New Mexico, Nevada, Oregon and Washington. Also the following silver districts in 4 Ash Peak, of the states mentioned: Vulture and Helvetia, Ariz: Miomi, Globe, Apache, Black Range, Chloride Flat, Georgetown and Lake Valley, New Mexico: Ashwood and Granite, oregon; Deertrail, Nespelam and Ruby-Conconully, Washington.

A. V. Heyl, Jr. - All districts east of the Mississippi River (except White Pine,

Harry Klemic and W. L. Newman - Eilver districts not associated with lead or zinc, in Arizona, New Mexico, Nevaña, Oregon, and Washington (except as listed above).

Size categories of deposits (as penciled in left margins)

	0	1	// 2	3
Cu	Less than 1,000 tons	1,000 to 50,000 tons	50,000 to 1,000,000 tons	More than 1,000,000 tons
Pb	11	#	11	<b>t</b> t
Zn	į į	Ħ	BE	Ħ
Λg	Less than 100,000 oz.	100,000 to 5,000,000 oz.	7 *	More than 1,000,000 oz.
Au	Less than 10,000 oz.	10,000 to 100.000 ox.	100,000 to 1,000,000 oz.	More than 1,000,000 oz.

(NOTE: Categories for Au are less certain than for others.)

District No. on metallogenie map peniled at lower right