

4960 0006

S27-28, 13N, 42E

New Moose

(256)

(Nye Co.)

109

Item 6

The New Moose tungsten property consists of 3 unpatented lode claims, known as the Moose group 1 to 3 situated on the east flank of the Toiyabe Range, 38 miles south of Austin and 136 miles south of Battle Mountain the shipping point on the Southern and Western Pacific Railroads.

Rocks exposed in the area consist of quartzites, schist, phyllite, and limestone that strike north and dip at low angles west. These formations have been broken by a north trending fault. Near the base of the limestone member, the limestone has been altered to tactite in zone 1,000 feet long and 1 to 3 feet thick.

In this altered zone, scheelite with varying small amounts of lead and silver occurs in widely spaced shoots.

Assay results of samples taken from the better mineralized sections indicate an ore grade of 0.4 percent WO_3 . No production or shipment of tungsten are reported from the property.

Silver Ace

USBM Unpubl. data, 1963

The Silver Ace property consist of 6 unpatented lode claims situated in a narrow westward trending canyon on the west flank of Bear Mountain, about 13 miles southeast of Beatty, and 83 miles northwest of Las Vegas, the supply and shipping point on the Union Pacific Railroad.

Two claims were originally located in 1929. From the work done at that time, about 40 tons of 2 ounce gold ore was produced.

The Silver Ace claim on which tungsten was discovered was located in 1933. From this claim 2 truck loads averaging 10.0 percent WO_3 were shipped in 1937. In February 1954, the property was optioned to

New Moose256
Item 6

The New Moose tungsten property consists of 3 unpatented lode claims, known as the Moose group 1 to 3 situated on the east flank of the Toyabe Range, 38 miles south of Austin and 136 miles south of Battle Mountain the shipping point on the Southern and Western Pacific Railroads.

Rocks exposed in the area consist of quartzites, schist, phyllite, and limestone that strike north and dip at low angles west. These formations have been broken by a north trending fault. Near the base of the limestone member, the limestone has been altered to tactite in zone 1,000 feet long and 1 to 3 feet thick.

In this altered zone, scheelite with varying small amounts of lead and silver occurs in widely spaced shoots.

Assay results of samples taken from the better mineralized sections indicate an ore grade of 0.4 percent WO_3 . No production or shipment of tungsten are reported from the property.